

CIDETRAK[®] IMM

A Mating Disruption Formulation for:

Indian meal moth (*Plodia interpunctella*), tobacco moth (*Ephestia elutella*), raisin moth (*Cadra figulilella*), almond moth (*Cadra cautella*), and Mediterranean flour moth (*Ephestia kuehniella*)

ACTIVE INGREDIENT:

(9Z,12E)-9,12-tetradecadien-1-yl acetate 3.2 %

OTHER INGREDIENTS: 96.8 %

TOTAL: 100.00%

NET CONTENTS:

Dispensers per package: 36 Units

One dispenser contains: 160 mgs, active ingredient

EPA Registration No. 51934-9

EPA Establishment No. 51934-OK-002

Batch Code _____

MANUFACTURED BY: TRÉCÉ INCORPORATED, 7569 Hwy 28 W · Adair, Oklahoma 74330

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse.

FIRST AID STATEMENT

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call poison control center or doctor for treatment advice.

HOT LINE NUMBER: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact Trécé, Inc. at 1-866-785-1313 between 8am and 6pm CST for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USING

CIDETRAK[®] IMM disrupts the mating communication between adult male and female moths, reduces fertile egg laying, suppresses subsequent larval infestation, and reduces the need for applying insecticides.

For maximum effectiveness, remove and destroy infested commodities and clean infested areas. Continuously monitor their facilities and commodities and maintain accurate records of target moth populations and larval damage within and near their facilities. Consult your local extension specialist, food safety consultant, pest management specialist, or Trece representative for assistance in designing the best CIDETRAK[®] IMM program for your commodities and facilities.

Facilities and commodities	Apply CIDETRAK [®] IMM in and around storage and manufacturing facilities containing raw grains and bean seeds and products including but not limited to wheat, barley, rye, oats, corn, popcorn, rice, all edible bean varieties, soybeans, cocoa beans; tobacco; birdseed; all spices; nuts including but not limited to peanuts, walnuts, almonds, pecans, hazelnuts, Brazil nuts and pistachios; dried commodities including but not limited to all fruits, vegetables, spices, nuts, and herbs, all animal and/or pet food, such as dog and cat food, cattle, hog, sheep and other animals in and around storage and manufacturing facilities including but not limited to raw, intermediate/in-process goods and finished goods facilities; mills, bakeries, drying areas.
Targeted Pests	Indian meal moth (<i>Plodia interpunctella</i>), Mediterranean flour moth (<i>Ephestia kuehniella</i>), tobacco moth (<i>Ephestia elutella</i>), raisin moth (<i>Cadra figulilella</i>), and almond moth (<i>Cadra cautella</i>).
Rate	Apply 1 dispenser per 3,531 to 10,594 Cu.Ft. (100 to 300 Cu. Meters) or 160 mgs. a.i. respectively depending on the infestation level. Apply 1 dispenser per 7,063 Cu.Ft. (200 Cu. Meters) to areas of moderate infestation. Apply the maximum dispenser rate to all buildings that are heavily infested with one or more target species, or adjacent to an untreated building, where access is evident. Do not exceed 200 grams of active ingredient per 7,063 Cu. Ft. (200 Cu. Meters) year.
Application	Place dispensers at a height of 5 to 16 feet (1.5 to 4.8 Meters) in a grid pattern over each floor of treated buildings. Attach dispensers securely to walls, pallets, columns, or ceiling hangers.
Timing	Apply prior to moth emergence in early spring or between any generation, in year round programs. Monitor moth activity using STORGARD II, PHEROCON IC traps and lures or STORGARD IMM oviposition traps. Begin monitoring in early spring and continue throughout the season to assess treatment effectiveness. Depending on temperature, CIDETRAK [®] IMM dispensers will release pheromone for 180 days. In areas with long field season, make a second application based on monitoring results. Note – CIDETRAK [®] IMM will not prevent stored commodity damage from immigration of mated female moths into treated buildings. Nearby (500 yards) untreated buildings can be a source of these mated females. Treatment with pheromone is recommended only when all buildings within or near treated buildings have been treated with CIDETRAK [®] IMM.
Treatment Tips	CIDETRAK [®] IMM suppresses mating of target insects. Immigration of mated female moths from adjacent, infested buildings can reduce the level of control. Area-wide treatment of entire sets of related buildings is the most effective strategy. To supplement CIDETRAK [®] IMM, use insecticides to control high populations. Monitor all pest populations to determine timely use of insecticides. To manage immigration and high population pressures, consider: 1. Treatment of external sources of infestation with CIDETRAK [®] IMM 2. Treatment of external sources of infestation with an insecticide. 3. Treatment of pheromone treated buildings with insecticide. Consult your local extension specialist, certified consultant, or Trece representative for local management strategies.

STORAGE AND DISPOSAL

Nonrefillable container. Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store unopened original package in a dry location at a temperature below 40° F.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Dispose of empty containers and foil envelopes in trash. Offer for recycling, if available.

WARRANTY AND LIMITATION OF DAMAGES

To the extent consistent with applicable law, all statements concerning the use of this product apply only when used as directed as stated on this product label. Trece Inc. warrants that the product conforms to this label and under normal conditions of use according to this label, is reasonably fit for the stated label purpose. The manufacturer makes no express or implied warranties of merchantability or fitness for a purpose concerning this product, or its use, extending beyond the label. Under no circumstance will Trece Inc. be liable for damage (special, consequential or incidental) that result from the handling, storage, or use of this product which is not in compliance with the label. Read all label directions carefully.

Buyer's exclusive remedy, should this product be defective, is replacement of the product, or if this is impractical, a purchase price refund.



Mailing Address: P.O. Box 129 · Adair, Oklahoma 74330
Shipping Address: 7569 Hwy 28 W. · Adair, Oklahoma 74330
Phone: (918) 785-3061 · Fax: (918) 785-3063

PHEROCON[®] and CIDETRAK[®] are registered Trademarks of TRECE Incorporated, Adair, OK USA
©2016 TRECE, Adair, OK USA · TRE0879